



* * * * *

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

* * * * *

Applicants: Sylvain RAULT, et al.
Serial No.: 10/689,394
Filed : October 20, 2003
Title : Imidazoline compounds.

* * * * *

HON. COMMISSIONER OF PATENTS AND TRADEMARKS
WASHINGTON, D.C. 20231

CLAIM TO PRIORITY AND FILING OF PRIORITY DOCUMENT
UNDER 37 CFR § 1.55 AND 35 USC § 119

Sir:

Herewith please find a certified copy of French priority application Serial No. 0213194 filed October 23, 2002, and certified translation thereof into English, the right of priority of which was claimed upon filing of the above-identified application, and which claim is hereby repeated.

Respectfully submitted,
THE FIRM OF HUESCHEN & SAGE

By: 
G. PATRICK SAGE, ATTORNEY

Dated: March 11, 2004.

Customer No. 25,666
500 Columbia Plaza
350 East Michigan Ave.
Kalamazoo, MI 49007
(616) 382-0030

Enclosures: Certified copy of French priority application Serial No. 0213194,
Certified translation thereof into English, and
Return postal card receipt.

CERTIFICATE OF MAILING UNDER 37 CFR 1.8(a)

I hereby certify that this correspondence is being deposited with the United States Postal Service as first-class mail in an envelope addressed to Commissioner of Patents and Trademarks, Alexandria, VA 22313-1450, on this date.

HUESCHEN AND SAGE


Dated: March 11, 2004



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

I, VIVIEN IRENE COULSON, declare:

1. That I am a citizen of the United Kingdom of Great Britain and Northern Ireland, residing at 96 Langley Road, Watford, Hertfordshire, WD17 4PJ;
2. That I am well acquainted with the French and English languages;
3. That the attached is a true translation into the English language of the certified copy of French Patent Application No. 0213194 filed 23 October 2002;
4. That I believe that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that wilful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such wilful false statements may jeopardise the validity of the patent application in the United States of America or any patent issuing thereon.

Declared this

15th

day of

November 2003

V.I. Coulson

V.I. COULSON

FRENCH REPUBLIC
NATIONAL INSTITUTE FOR INDUSTRIAL PROPERTY

PATENT OF INVENTION

UTILITY CERTIFICATE - CERTIFICATE OF ADDITION

OFFICIAL COPY

The Director General of the National Institute for Industrial Property certifies that the attached document is the true certified copy of an application for an Industrial Property Right filed at the Institute.

Issued in Paris, 20 JUNE 2003

For the Director General of the
National Institute for
Industrial Property
The Head of the Patents Division

(signature)

Martine PLANCHE

NATIONAL INSTITUTE FOR INDUSTRIAL PROPERTY

HEAD OFFICE

26 bis, rue de Saint Petersburg
75800 PARIS cedex 08

Tel.: 33 (0)1 53 04 53 04 Fax: 33 (0)1 53 04 45 23 www.inpi.fr

NATIONAL PUBLIC INSTITUTION CREATED BY LAW NO. 51-444 of 19 April 1951

**INPI NATIONAL INSTITUTE
FOR INDUSTRIAL PROPERTY**

26 bis, rue de Saint Pétersbourg
75800 Paris Cedex 08
Tel.: 01 53 04 53 04

Fax: 01 42 94 86 54

**PATENT OF INVENTION
UTILITY CERTIFICATE**

Intellectual Property Code - Book VI

REQUEST FOR GRANT 1/2

Reserved for INPI

This form is to be completed legibly in black ink

DEPOSITION OF DOCUMENTS DATE 23 OCT 2002 PLACE 75 INPI PARIS NATIONAL REGISTRATION NO. ATTRIBUTED BY THE INPI 0213194 FILING DATE ATTRIBUTED BY THE INPI 23 OCT. 2002		1 NAME AND ADDRESS OF THE APPLICANT OR OF THE AUTHORISED AGENT TO WHOM CORRESPONDENCE MUST BE ADDRESSED LES LABORATOIRES SERVIER Direction Brevets 12, Place de La Défense 92415 COURBEVOIE Cedex FRANCE	
Your references for this file (optional) 70097			
Confirmation of a deposit by facsimile		<input type="checkbox"/> No. attributed by the INPI to the facsimile	
2 NATURE OF THE APPLICATION		Mark one of the following 4 boxes	
Patent Application		<input checked="" type="checkbox"/>	
Application for a Utility Certificate		<input type="checkbox"/>	
Divisional Application <i>Initial patent application</i> <i>or initial utility certificate application</i>		<input type="checkbox"/> No. Date No. Date	
Conversion of a European Patent Application <i>Initial patent application</i>		<input type="checkbox"/> No. Date	
3 TITLE OF THE INVENTION (maximum 200 characters or spaces) New imidazoline compounds, a process for their preparation and pharmaceutical compositions containing them			
4 DECLARATION OF PRIORITY OR REQUEST FOR THE BENEFIT OF THE FILING DATE OF A PRIOR FRENCH APPLICATION		Country or organisation Date No. Country or organisation Date No. Country or organisation Date No. <input type="checkbox"/> If there are other priorities, mark the box and use the "Continuation" form	
5 APPLICANT		<input type="checkbox"/> If there are other Applicants, mark the box and use the "Continuation" form	
Name or company name		LES LABORATOIRES SERVIER	
Forenames			
Legal nature			
SIREN No.			
APE-NAF Code			
Address	Street	12, Place de La Défense	
	Postal code and town	92415	COURBEVOIE Cedex
Country		FRANCE	
Nationality		FRENCH	
Telephone no. (optional)		01.55.72.60.00	
Fax no. (optional)		01.55.72.72.13	
E-mail address (optional)			

**INPI NATIONAL INSTITUTE
FOR INDUSTRIAL PROPERTY**
**PATENT OF INVENTION
UTILITY CERTIFICATE**

REQUEST FOR GRANT 2/2

Reserved for INPI


DEPOSITION OF DOCUMENTS			
DATE 23 OCT 2002			
PLACE 75 INPI PARIS			
NATIONAL REGISTRATION NO. ATTRIBUTED BY THE INPI 0213194			
Your references for this file: (optional)		70097	
6 AUTHORISED AGENT			
Name		RIVIERE	
Forename		François	
Practice or company		LES LABORATOIRES SERVIER	
No. of standing power of attorney and/or of contractual bond			
Address	Street	12, Place de La Défense	
	Postal code and town	92415	COURBEVOIE Cedex
Telephone no. (optional)		01.55.72.60 00	
Fax no. (optional)		01.55.72.72.13	
E-mail address (optional)			
7 INVENTOR(S)			
The inventors are the Applicants		<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No In this case, supply a separate declaration of inventorship	
8 SEARCH REPORT		For a patent application only (including division and conversion)	
immediate drawing up or deferred drawing up		<input checked="" type="checkbox"/> <input type="checkbox"/>	
Payment of the fees in instalments		Payment in three instalments, for natural persons only <input type="checkbox"/> Yes <input type="checkbox"/> No	
9 REDUCTION IN FEES		For natural persons only <input type="checkbox"/> Requested for the first time for this invention (attach a notice of non-imposition) <input type="checkbox"/> Requested prior to this deposit (attach a copy of the admissibility decision for this invention or indicate its reference)	
If you have used the "Continuation" form, indicate the number of pages attached			
10 SIGNATURE OF THE APPLICANT OR OF THE AUTHORISED AGENT (Name and position of signatory)		STAMP OF THE PREFECTURE OR OF THE INPI	
François RIVIERE, Patent Engineer		(signature) L. MARIELLO	

The invention relates to new imidazoline compounds, to a process for their preparation and to pharmaceutical compositions containing them.

As far as chemical structures are concerned, the literature provides numerous examples of imidazoline compounds. They have been described, *inter alia*, for their therapeutic use. Imidazoline compounds are known, for example, to have cardiotonic properties (GB 119/963), α -adrenergic stimulating properties (Eur. J. Med. Chem., 1989, 24(6), 619, J. Pharmacobio. Dyn., 1986, 9(4), 395), antidepressant and anti-inflammatory properties (US 3932-431), antihyperglycaemic and antidiabetic properties (EP 924 209, EP 1 145 717, EP 288 978, JP 04178381 and WO 02 38559) or to be capable of treating pathologies associated with imidazoline receptors (EP 846 688).

The Applicant has now discovered new imidazoline compounds having a novel cycloalkylimidazoline structure that gives them antidiabetic properties without side effects owing to the absence of serotonin reuptake inhibition.

The present invention relates more especially to compounds of formula (I) :



wherein :

- R^1 represents an optionally substituted heteroaryl group,
- R^2 represents an optionally substituted cycloalkyl group,
- R^3 represents a hydrogen atom or an alkyl group, and
- R^4 and R^5 , which may be identical or different, each represents a hydrogen atom, a halogen atom or an alkyl, polyhaloalkyl, $R^{10}-C(X)-R^{11}-$, $R^{10}-Y-C(X)-R^{11}-$, $R^{10}-C(X)-Y-R^{11}-$, $R^{10}-Y-R^{11}-$

or $R^{10}-S(O)_n-R^{11}$ group,

in which :

- R^{10} represents a hydrogen atom or an alkyl group,
- R^{11} represents a bond, or an alkylene, alkenylene or alkynylene group,
- 5 - X represents an oxygen atom, a sulphur atom, or an NR^{12} group in which R^{12} represents a hydrogen atom or an alkyl group,
- Y represents an oxygen atom, a sulphur atom, or an amino or alkylamino group, and
- n represents an integer of from 1 to 2 inclusive,

10 to their enantiomers, diastereoisomers, tautomers, and also addition salts thereof with a pharmaceutically acceptable acid or base,

it being understood that:

- the term "alkyl" denotes a linear or branched hydrocarbon chain containing from 1 to 6 carbon atoms,
- the term "alkoxy" denotes an alkyl-oxy group in which the alkyl chain, which may be
15 linear or branched, contains from 1 to 6 carbon atoms,
- the term "alkylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms,
- the term "alkenylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms and from 1 to 3 double bonds,
- the term "alkynylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms and from 1 to 3 triple bonds,
- 20 - the term "polyhaloalkyl" denotes a linear or branched carbon chain containing from 1 to 3 carbon atoms and from 1 to 7 halogen atoms,
- the term "heteroaryl" denotes a mono- or bi-cyclic group having from 5 to 11 ring members in which at least one of the rings is aromatic and containing in the monocycle or in the bicycle 2 or 3 hetero atoms selected from nitrogen, oxygen and sulphur, and
- 25 - the term "cycloalkyl" denotes a hydrocarbon monocycle or bicycle that contains from 3 to 10 carbon atoms and is optionally unsaturated by 1 or 2 unsaturated bonds;

or $R^{10}-S(O)_n-R^{11}$ group,

in which :

- R^{10} represents a hydrogen atom or an alkyl group,
- R^{11} represents a bond, or an alkylene, alkenylene or alkynylene group,
- 5 - X represents an oxygen atom, a sulphur atom, or an NR^{12} group in which R^{12} represents a hydrogen atom or an alkyl group,
- Y represents an oxygen atom, a sulphur atom, or an amino or alkylamino group, and
- n represents an integer of from 1 to 2 inclusive,

10 to their enantiomers, diastereoisomers, tautomers, and also addition salts thereof with a pharmaceutically acceptable acid or base,

it being understood that:

- the term "alkyl" denotes a linear or branched hydrocarbon chain containing from 1 to 6 carbon atoms,
- the term "alkoxy" denotes an alkyl-oxy group in which the alkyl chain, which may be
15 linear or branched, contains from 1 to 6 carbon atoms,
- the term "alkylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms,
- the term "alkenylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms and from 1 to 3 double bonds,
- the term "alkynylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms and from 1 to 3 triple bonds,
- 20 - the term "polyhaloalkyl" denotes a linear or branched carbon chain containing from 1 to 3 carbon atoms and from 1 to 7 halogen atoms,
- the term "heteroaryl" denotes a mono- or bi-cyclic group having from 5 to 11 ring members in which at least one of the rings is aromatic and containing in the monocycle or in the bicycle 1, 2 or 3 hetero atoms selected from nitrogen, oxygen and sulphur, and
- 25 - the term "cycloalkyl" denotes a hydrocarbon monocycle or bicycle that contains from 3 to 10 carbon atoms and is optionally unsaturated by 1 or 2 unsaturated bonds;

- 5 - the expression "optionally substituted" associated with the terms cycloalkyl and heteroaryl denotes that the groups in question are substituted by one or two identical or different substituents selected from halogen atoms and the groups alkyl, alkoxy, hydroxy, cyano, nitro, amino (optionally substituted by one or two alkyl groups) and -C(O)R_d wherein R_d represents a group selected from hydroxy, alkoxy and amino, it being understood that the heteroaryl group may be additionally substituted by an oxo group on the non-aromatic moiety of the heteroaryl.

10 Among the pharmaceutically acceptable acids there may be mentioned, without implying any limitation, hydrochloric, hydrobromic, sulphuric, phosphonic, acetic, trifluoroacetic, lactic, pyruvic, malonic, succinic, glutaric, fumaric, tartaric, maleic, citric, ascorbic, methanesulphonic and camphoric acid etc...

Among the pharmaceutically acceptable bases there may be mentioned, without implying any limitation, sodium hydroxide, potassium hydroxide, triethylamine, etc...

15 An advantageous aspect of the invention concerns compounds wherein R⁴ and R⁵, which may be identical or different, each represents a hydrogen atom or an alkyl group.

Another advantageous aspect concerns compounds of formula (I) wherein R³ represents a hydrogen atom.

The preferred alkyl group of the invention for the groups R⁴ and/or R⁵ is the methyl group.

20 Preferred cycloalkyl groups of the invention are the groups cyclopentyl, cyclohexyl and cycloheptyl, and more especially the cyclohexyl group.

Preferred heteroaryl groups of the invention are aromatic groups having 5 or 6 ring members, such as furyl, thienyl, pyrrolyl and pyridyl and, more especially, thienyl.

- 5 - the expression "optionally substituted" associated with the terms cycloalkyl and heteroaryl denotes that the groups in question are substituted by one or two identical or different substituents selected from halogen atoms and the groups alkyl, alkoxy, hydroxy, cyano, nitro, amino (optionally substituted by one or two alkyl groups) and -C(O)R_d wherein R_d represents a group selected from hydroxy, alkoxy and amino, it being understood that the heteroaryl group may be additionally substituted by an oxo group on the non-aromatic moiety of the heteroaryl.

10 Among the pharmaceutically acceptable acids there may be mentioned, without implying any limitation, hydrochloric, hydrobromic, sulphuric, phosphonic, acetic, trifluoroacetic, lactic, pyruvic, malonic, succinic, glutaric, fumaric, tartaric, maleic, citric, ascorbic, methanesulphonic and camphoric acid etc...

Among the pharmaceutically acceptable bases there may be mentioned, without implying any limitation, sodium hydroxide, potassium hydroxide, triethylamine, etc...

15 An advantageous aspect of the invention concerns compounds wherein R⁴ and R⁵, which may be identical or different, each represents a hydrogen atom or an alkyl group.

Another advantageous aspect concerns compounds of formula (I) wherein R³ represents a hydrogen atom.

The preferred alkyl group of the invention for the groups R⁴ and/or R⁵ is the methyl group.


20 Preferred R² groups of the invention are the groups cyclopentyl, cyclohexyl and cycloheptyl, optionally substituted by an alkyl group, and more especially the cyclohexyl group.

Preferred heteroaryl groups of the invention are aromatic groups having 5 or 6 ring members, such as furyl, thienyl, pyrrolyl and pyridyl and, more especially, thienyl.

An especially advantageous aspect of the invention concerns compounds of formula (I) wherein R^1 represents a heteroaryl group having 5 or 6 optionally substituted ring members, R^2 represents a cyclohexyl group optionally substituted by an alkyl group, R^3 represents a hydrogen atom, and R^4 and R^5 , which may be identical or different, each represents a hydrogen atom or an alkyl group.


Among the preferred compounds of the invention, the following may be mentioned more especially: 2-[cyclohexyl(3-thienyl)methyl]-4-methyl-4,5-dihydro-1*H*-imidazole and its tautomer 2-[cyclohexyl(3-thienyl)methyl]-5-methyl-4,5-dihydro-1*H*-imidazole.

The invention relates also to a process for the preparation of compounds of formula (I), which is characterised in that there is used as starting material a compound of formula (II) :



wherein R^1 , R^2 and R^3 are as defined for formula (I),

which compounds of formula (II) are condensed with a diamine (III) :



wherein R^4 and R^5 are as defined for formula (I)

to yield, in the presence of an appropriate catalyst, compounds of formula (I),


- which may optionally be purified according to a conventional purification technique,
- which, if desired, are separated into their stereoisomers according to a conventional separation technique,
- which, if desired, are converted into addition salts with a pharmaceutically acceptable

-4 -

An especially advantageous aspect of the invention concerns compounds of formula (I) wherein R^1 represents a heteroaryl group having 5 or 6 optionally substituted ring members, R^2 represents a cyclohexyl or cycloheptyl group optionally substituted by an alkyl group, R^3 represents a hydrogen atom, and R^4 and R^5 , which may be identical or different, each represents a hydrogen atom or an alkyl group.


Among the preferred compounds of the invention, the following may be mentioned more especially: 2-[cyclohexyl(3-thienyl)methyl]-4-methyl-4,5-dihydro-1*H*-imidazole and its tautomer 2-[cyclohexyl(3-thienyl)methyl]-5-methyl-4,5-dihydro-1*H*-imidazole.

The invention relates also to a process for the preparation of compounds of formula (I), which is characterised in that there is used as starting material a compound of formula (II) :



wherein R^1 , R^2 and R^3 are as defined for formula (I),

which compounds of formula (II) are condensed with a diamine (III) :



wherein R^4 and R^5 are as defined for formula (I)

to yield, in the presence of an appropriate catalyst, compounds of formula (I),

- which may optionally be purified according to a conventional purification technique,
- which, if desired, are separated into their stereoisomers according to a conventional separation technique,
- which, if desired, are converted into addition salts with a pharmaceutically acceptable it

acid or base,

it being understood that

- at any moment considered appropriate during the course of the process described above, the carbonyl, thiocarbonyl, amino or alkylamino group or groups of the starting reagents (II) and (III) may be protected and then, after condensation, deprotected for the purposes of the synthesis,
- the reagents (II) and (III) are prepared according to known procedures described in the literature.

The compounds exhibit, *inter alia*, excellent activity in reducing blood glucose levels. Those properties justify their therapeutic use in the treatment and/or prophylaxis of hyperglycaemia, dyslipidaemia and, more especially, in the treatment of non-insulin-dependent type II diabetes, obesity, glucose intolerance, and diabetic complications, especially in respect of the cardiovascular system.

The activity of the compounds is likewise recommended for the treatment and/or prophylaxis of other diseases, including type I diabetes, hypertriglyceridaemia, metabolic syndrome, insulin resistance, dyslipidaemia in diabetics, hyperlipidaemia and hypercholesterolaemia.

The compounds of the present invention in addition have a very weak affinity for serotonin receptor sites, and they are of low toxicity, unlike the prior art compounds.

The present invention relates also to pharmaceutical compositions comprising as active ingredient at least one compound of formula (I), on its own or in combination with one or
5 more pharmaceutically acceptable, inert, non-toxic excipients or carriers.

Among the pharmaceutical compositions according to the invention there may be mentioned more especially those which are suitable for oral, parenteral or nasal administration, tablets or dragées, sublingual tablets, gelatin capsules, lozenges, suppositories, creams, ointments, dermal gels, etc..

The useful dosage varies in accordance with the age and weight of the patient, the nature and the severity of the disorder, and also the administration route, which may be oral, nasal, rectal or parenteral. Generally, the unit dosage ranges from 0.1 to 500 mg for a treatment of from 1 to 3 administrations per 24 hours.

5 The following Examples illustrate the invention and do not limit it in any way. The structures of the described compounds were confirmed by customary spectroscopic and spectrometric techniques.

The starting materials used are known products or products prepared according to known procedures.

10 The name of the example compounds hereinbelow also includes the tautomers thereof, for example 2-[cyclohexyl(3-thienyl)methyl]-4-methyl-4,5-dihydro-1*H*-imidazole and its tautomer 2-[cyclohexyl(3-thienyl)methyl]-5-methyl-4,5-dihydro-1*H*-imidazole.

**GENERAL PROTOCOL A : synthesis of cycloalkylidene acetonitrile compounds
(PREPARATIONS 1 to 6) :**

15 0.1 mol of acetonitrile compound is added to a solution of 0.1 mol of sodium ethanolate in 50 ml of ethanol, and then 0.1 mol of ketone compound is added. The mixture is heated at 60°C for 4 hours, and the resulting solution is then concentrated under reduced pressure to half the initial volume and subsequently poured into 200 ml of water. Following extraction with ether, the ethereal fractions are washed with water and dried over magnesium sulphate
20 and the solvent is evaporated off. The product so obtained will be used in the following step without additional purification.

**GENERAL PROTOCOL B : reduction of the nitrogen-containing compounds
(PREPARATIONS 7 to 12) :**

0.125 mol of sodium borohydride is added to a solution of 0.05 mol of compounds of
25 Preparations 1-6 in 100 ml of ethanol. The solution is heated with reflux of ethanol for

24 hours and the ethanol is evaporated off under reduced pressure. The residue is taken up in 500 ml of water and extracted with ether. The desired product is obtained after drying the organic phase over magnesium sulphate and evaporating off the ether. The product so obtained will be used in the following step without additional purification.

5 **PREPARATION 1 : Cyclopentylidene(3-thienyl)acetonitrile**

The expected intermediate is obtained in the manner described in General Protocol A starting from 3-thienylacetonitrile and cyclopentanone.

● **PREPARATION 2 : Cyclohexylidene(2-thienyl)acetonitrile**

10 The expected intermediate is obtained in the manner described in General Protocol A starting from 2-thienylacetonitrile and cyclohexanone.

● **PREPARATION 3 : Cyclohexylidene(3-thienyl)acetonitrile**

The expected intermediate is obtained in the manner described in General Protocol A starting from 3-thienylacetonitrile and cyclohexanone.

● **PREPARATION 4 : (4-Methylcyclohexylidene)(3-thienyl)acetonitrile**

15 The expected intermediate is obtained in the manner described in General Protocol A starting from 3-thienylacetonitrile and 4-methylcyclohexanone.

● **PREPARATION 5 : Cycloheptalidene(3-thienyl)acetonitrile**

The expected intermediate is obtained in the manner described in General Protocol A starting from 3-thienylacetonitrile and cycloheptanone.

PREPARATION 6 : Cyclohexylidene(1-methyl-1*H*-pyrrol-2-yl)acetonitrile

The expected intermediate is obtained in the manner described in General Protocol A starting from (1-methyl-1*H*-pyrrol-3-yl)acetonitrile and cyclohexanone.

PREPARATION 7 : Cyclopentyl(3-thienyl)acetonitrile

- 5 The expected intermediate is obtained in the manner described in General Protocol B starting from the compound of Preparation 1.

PREPARATION 8 : Cyclohexyl(2-thienyl)acetonitrile

The expected intermediate is obtained in the manner described in General Protocol B starting from the compound of Preparation 2.

10 **PREPARATION 9 : Cyclohexyl(3-thienyl)acetonitrile**

The expected intermediate is obtained in the manner described in General Protocol B starting from the compound of Preparation 3.

PREPARATION 10 : (4-Methylcyclohexyl)(3-thienyl)acetonitrile

- 15 The expected intermediate is obtained in the manner described in General Protocol B starting from the compound of Preparation 4.

PREPARATION 11 : Cycloheptyl(3-thienyl)acetonitrile

The expected intermediate is obtained in the manner described in General Protocol B starting from the compound of Preparation 5.

PREPARATION 12 : Cyclohexyl(1-methyl-1*H*-pyrrol-2-yl)acetonitrile

The expected intermediate is obtained in the manner described in General Protocol B starting from the compound of Preparation 6.

PREPARATION 13 : Cyclohexyl(pyrid-2-yl)acetonitrile

5 0.1 mol of 2-pyridylacetonitrile in 10 ml of benzene is added to a suspension of 0.011 mol of sodium amide in 15 ml of anhydrous benzene. The reaction mixture is heated at reflux for 3 hours. The solution is then allowed to return to ambient temperature in order to enable the dropwise addition of 0.1 mol of bromocyclohexane, and then the solution is again heated at reflux for 8 hours.

10 The mixture is cooled and 50 ml of water are added to eliminate the excess of sodium amide. The organic phase is extracted and dried over magnesium sulphate and the solvent is evaporated off under reduced pressure.

GENERAL PROTOCOL C : synthesis of the compounds of EXAMPLES 1 to 12 :

15 A catalytic amount of phosphorus pentasulphide (0.5 g, 0.001 mol) is added to 0.02 mol of the compounds of Preparations 7 to 13 in 25 ml of the chosen diamine compound. The mixture is heated at reflux for 5 hours. The solution is then poured into 50 ml of water and extracted twice with 50 ml of dichloromethane. Following evaporation of the solvent, the product is recrystallised from a minimum amount of acetonitrile.

EXAMPLE 1 : 2-[Cyclopentyl(3-thienyl)methyl]-4,5-dihydro-1*H*-imidazole

20 The title compound is obtained according to General Protocol C starting from the compound of Preparation 7 and 1,2-ethanediamine.

Melting point : 157°C.

EXAMPLE 2 : 2-[Cyclohexyl(2-thienyl)methyl]-4,5-dihydro-1H-imidazole

The title compound is obtained according to General Protocol C starting from the compound of Preparation 8 and 1,2-ethanediamine.

Melting point : 155°C.

EXAMPLE 3 : 2-[Cyclohexyl(3-thienyl)methyl]-4,5-dihydro-1H-imidazole

The title compound is obtained according to General Protocol C starting from the compound of Preparation 9 and 1,2-ethanediamine.

Melting point : 182°C.

**EXAMPLE 4 : 2-[Cyclohexyl(3-thienyl)methyl]-4,5-dihydro-1H-imidazole,
enantiomer 1**

By separating the compound described above in Example 3 by chiral chromatography, one of the enantiomers is isolated.

**EXAMPLE 5 : 2-[Cyclohexyl(3-thienyl)methyl]-4,5-dihydro-1H-imidazole,
enantiomer 2**

By separating the compound described above in Example 3 by chiral chromatography, the other enantiomer is isolated.

EXAMPLE 6 : 2-[Cyclohexyl(3-thienyl)methyl]-4-methyl-4,5-dihydro-1H-imidazole

The title compound is obtained according to General Protocol C starting from the compound of Preparation 9 and 1,2-propanediamine.

Melting point : 155°C.

EXAMPLE 7 : 2-[Cyclohexyl(3-thienyl)methyl]-4,4-dimethyl-4,5-dihydro-1H-imidazole

The title compound is obtained according to General Protocol C starting from the compound of Preparation 9 and 2-methyl-1,2-propanediamine.

5 Melting point : 162°C.

EXAMPLE 8 : 2-[(4-Methylcyclohexyl)(3-thienyl)methyl]-4,5-dihydro-1H-imidazole

The title compound is obtained according to General Protocol C starting from the compound of Preparation 10 and 1,2-ethanediamine.

Melting point : 168°C.

10 **EXAMPLE 9 : 2-[Cyclohexyl(1-methyl-1H-pyrrol-2-yl)methyl]-4,5-dihydro-1H-imidazole**

The title compound is obtained according to General Protocol C starting from the compound of Preparation 12 and 1,2-ethanediamine.

Melting point : 120°C.

15 **EXAMPLE 10 : 2-[Cyclohexyl(4,5-dihydro-1H-imidazol-2-yl)methyl]pyridine**

The title compound is obtained according to General Protocol C starting from the compound of Preparation 13 and 1,2-ethanediamine.

Melting point : 116°C.

EXAMPLE 11 : 2-[Cycloheptyl(3-thienyl)methyl]-4,5-dihydro-1H-imidazole

20 The title compound is obtained according to General Protocol C starting from the compound of Preparation 11 and 1,2-ethanediamine.

Melting point : 155°C.

EXAMPLE 12 : 2-[Cycloheptyl(3-thienyl)methyl]-4-methyl-4,5-dihydro-1H-imidazole

The title compound is obtained according to General Protocol C starting from the compound of Preparation 11 and 1,2-propanediamine.

Melting point : 125°C.

5

PHARMACOLOGICAL STUDY**EXAMPLE A : Hypoglycaemic activity**

The hypoglycaemic activity of the compounds of the invention was examined in three-month-old male Wistar rats weighing about 250 g. An experimental diabetes is obtained by the *iv* injection, under ketamine hydrochloride anaesthesia, of a low dose of streptozotocin (35 mg/kg *iv*) dissolved in a citrate buffer. Such rats are called STZ rats and are characterised by a slight basal hyperglycaemia, a marked intolerance to glucose and a clear change in the secretion of insulin.

Homeostasis was evaluated by a glucose tolerance test, carried out two weeks after injection with streptozotocin.

Finally, the hypoglycaemic activity was evaluated in "Zucker" rats. "Zucker fatty" fa/fa rats arose following a spontaneous mutation of the 13M strain (Zucker & Zucker, 1961), and are genetically insulin-resistant and obese.

Their obesity can be observed from the age of four weeks, and is thus accompanied by insulin resistance, hyperinsulinaemia and hyperlipidaemia. This model is predictive of diabetic conditions exhibiting associated metabolic disorders, such as obesity.

Homeostasis was evaluated likewise by a glucose tolerance test.

. Oral glucose tolerance test (OGTT)

Glucose is administered *per os* (2 g/kg) to conscious rats. Blood samples are collected before and 10, 20, 30, 40, 60, 90 and 120 minutes after the glucose administration.

The product to be tested is administered *per os* 1 hour before the OGTT, and the control animals receive solvent (gum arabic).

The compounds of the invention reduce glycaemia very significantly.

For example, at 10 mg/kg, the compound of Example 6 reduces glycaemia by 13 %, 18 %
5 and 14 % in non-diabetic Wistar rats, STZ Wistar rats and Zucker rats, respectively.

EXAMPLE B : Hypolipaeic activity

The products of the invention were tested *in vivo* in the obese ob/ob mouse, used as a model of obesity-associated insulin resistance. By way of example, the compound of Example 6 significantly lowers the triglycerides (after chronic administration by the oral
10 route at a dose of 30 mg/kg/day for 4 days) by 32 % (animals treated with the compound of Example 6 vs. untreated animals).

In this model, the compounds of the invention are thus shown also to be powerful hypolipaeics.

EXAMPLE C : Determination of the affinity for serotonin reuptake sites in the rat

15 The affinity of the compounds of the invention was determined by competition experiments with [³H]-paroxetine. The membranes are prepared from the rat frontal cortex and incubated in triplicate for 2 hours, at 25°C, with 0.25 nM [³H]-paroxetine and the cold ligand in a final volume of 0.4 ml. The incubation buffer contains 50 mM TRIS-HCl (pH 7.4), 120 mM NaCl and 5 mM KCl. The non-specific binding is determined using 10 µM
20 citalopram. At the end of incubation, the medium is filtered through filters and washed three times with 5 ml of cooled buffer. The radioactivity retained on the filters is determined by liquid scintillation counting. The binding isotherms are analysed by non-linear regression in order to determine the IC₅₀ values.

It appears that the compounds of the invention exhibit a weaker affinity for the serotonin reuptake sites than the compounds of the prior art, and thus a decrease in central toxicity confirmed by the Irwin test (see Example D).

- 5 By way of example, the compound of Example 3 has an IC_{50} of $2 \times 10^{-6} M$, which is superior to that of the prior art (EP 846 688) : 2-[cyclohexyl(phenyl)methyl]-4,5-dihydro-1*H*-imidazole ($1.5 \times 10^{-7} M$).


EXAMPLE D : Acute toxicity study – Irwin Test

10 Three rats per dose are treated *per os* with one of the compounds of the invention (dispersed in 0.5 % carboxymethyl cellulose in distilled water) and are observed at regular intervals after 24 hours. The presence or absence of the following symptoms are recorded : mortality, sedation, excitation, aggressiveness, tail form, convulsions, pain, trembling, exophthalmia, salivation, piloerection, defecation, fear, etc, according to the criteria described by Irwin (Psychopharmacologia, 1968, 13, 222). This test enables the toxicity
15 and the effect on behaviour to be evaluated.

It appears that the compounds of the invention, according to the therapeutic index (ratio of the minimum active dose to the dose at which symptoms appear in the central nervous system), are less toxic than those of the prior art.

EXAMPLE E : Pharmaceutical composition

20 Formulation for the preparation of 1000 tablets each containing a dose of 10 mg :
compound of Example 6 10 g
hydroxypropyl cellulose 2 g
wheat starch 10 g
lactose 100 g
25 magnesium stearate..... 3 g
talc..... 3 g

CLAIMS**1. Compounds of formula (I) :**

wherein :

- R¹ represents an optionally substituted heteroaryl group,
- R² represents an optionally substituted cycloalkyl group,
- R³ represents a hydrogen atom or an alkyl group, and
- R⁴ and R⁵, which may be identical or different, each represents a hydrogen atom, a halogen atom or an alkyl, polyhaloalkyl, R¹⁰—C(X)—R¹¹—, R¹⁰—Y—C(X)—R¹¹—, R¹⁰—C(X)—Y—R¹¹—, R¹⁰—Y—R¹¹— or R¹⁰—S(O)_n—R¹¹— group,

in which :

- R¹⁰ represents a hydrogen atom or an alkyl group,
- R¹¹ represents a bond, or an alkylene, alkenylene or alkynylene group,
- X represents an oxygen atom, a sulphur atom, or an NR¹² group in which R¹² represents a hydrogen atom or an alkyl group,
- Y represents an oxygen atom, a sulphur atom, or an amino or alkylamino group, and
- n represents an integer of from 1 to 2 inclusive,

the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base,

it being understood that:

- the term "alkyl" denotes a linear or branched hydrocarbon chain containing from 1 to 6 carbon atoms,
- the term "alkoxy" denotes an alkyl-oxy group in which the alkyl chain, which may be linear or branched, contains from 1 to 6 carbon atoms,
- the term "alkylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms,
- the term "alkenylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms and from 1 to 3 double bonds,
- the term "alkynylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms and from 1 to 3 triple bonds,
- the term "polyhaloalkyl" denotes a linear or branched carbon chain containing from 1 to 3 carbon atoms and from 1 to 7 halogen atoms,
- the term "heteroaryl" denotes a mono- or bi-cyclic group having from 5 to 11 ring members in which at least one of the rings is aromatic and containing in the monocycle or in the bicycle 2 or 3 hetero atoms selected from nitrogen, oxygen and sulphur, and
- the term "cycloalkyl" denotes a hydrocarbon monocycle or bicycle that contains from 3 to 10 carbon atoms and is optionally unsaturated by 1 or 2 unsaturated bonds;
- the expression "optionally substituted" associated with the terms cycloalkyl and heteroaryl denotes that the groups in question are substituted by one or two identical or different substituents selected from halogen atoms and the groups alkyl, alkoxy, hydroxy, cyano, nitro, amino (optionally substituted by one or two alkyl groups) and -C(O)R_d wherein R_d represents a group selected from hydroxy, alkoxy and amino, it being understood that the heteroaryl group may be additionally substituted by an oxo group on the non-aromatic moiety of the heteroaryl.

2. Compound of formula (I) according to claim 1 wherein R⁴ and R⁵, which may be identical or different, each represents a hydrogen atom or an alkyl group, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.

being understood that:

- the term "alkyl" denotes a linear or branched hydrocarbon chain containing from 1 to 6 carbon atoms,
- the term "alkoxy" denotes an alkyl-oxy group in which the alkyl chain, which may be linear or branched, contains from 1 to 6 carbon atoms,
- the term "alkylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms,
- the term "alkenylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms and from 1 to 3 double bonds,
- the term "alkynylene" denotes a linear or branched bivalent hydrocarbon chain containing from 1 to 6 carbon atoms and from 1 to 3 triple bonds,
- the term "polyhaloalkyl" denotes a linear or branched carbon chain containing from 1 to 3 carbon atoms and from 1 to 7 halogen atoms,
- the term "heteroaryl" denotes a mono- or bi-cyclic group having from 5 to 11 ring members in which at least one of the rings is aromatic and containing in the monocycle or in the bicycle 1, 2 or 3 hetero atoms selected from nitrogen, oxygen and sulphur, and
- the term "cycloalkyl" denotes a hydrocarbon monocycle or bicycle that contains from 3 to 10 carbon atoms and is optionally unsaturated by 1 or 2 unsaturated bonds;
- the expression "optionally substituted" associated with the terms cycloalkyl and heteroaryl denotes that the groups in question are substituted by one or two identical or different substituents selected from halogen atoms and the groups alkyl, alkoxy, hydroxy, cyano, nitro, amino (optionally substituted by one or two alkyl groups) and -C(O)R_d wherein R_d represents a group selected from hydroxy, alkoxy and amino, it being understood that the heteroaryl group may be additionally substituted by an oxo group on the non-aromatic moiety of the heteroaryl.

2. Compound of formula (I) according to claim 1 wherein R⁴ and R⁵, which may be identical or different, each represents a hydrogen atom or an alkyl group, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.

3. Compound of formula (I) according to either claim 1 or claim 2 wherein R^3 represents a hydrogen atom, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.
- 5 4. Compound of formula (I) according to any one of claims 1 to 3 wherein R^1 represents an optionally substituted heteroaryl group having 5 or 6 ring members, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.
- 10 5. Compound of formula (I) according to any one of claims 1 to 4 wherein R^2 represents a cycloalkyl group having 5, 6 or 7 ring members, which is optionally substituted by an alkyl group, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.
- 15 6. Compound of formula (I) according to any one of claims 1 to 5 wherein R^1 represents an optionally substituted heteroaryl group having 5 or 6 ring members, R^2 represents a cyclohexyl or cycloheptyl group optionally substituted by an alkyl group, R^3 represents a hydrogen atom and R^4 and R^5 , which may be identical or different, each represents a hydrogen atom or an alkyl group, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.
- 20 7. Compound of formula (I) according to any one of claims 1, 2, 5 and 6 wherein the alkyl group is a methyl group, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.
- 25 8. Compound of formula (I) according to any one of claims 1 to 7 which is 2-[cyclohexyl(3-thienyl)methyl]-4-methyl-4,5-dihydro-1*H*-imidazole, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid.

3. Compound of formula (I) according to either claim 1 or claim 2 wherein R^3 represents a hydrogen atom, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.

5 4. Compound of formula (I) according to any one of claims 1 to 3 wherein R^1 represents an optionally substituted heteroaryl group having 5 or 6 ring members, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.


10 5. Compound of formula (I) according to any one of claims 1 to 4 wherein R^2 represents a cyclopentyl, cyclohexyl or cycloheptyl group optionally substituted by an alkyl group, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.

15 6. Compound of formula (I) according to any one of claims 1 to 5 wherein R^1 represents an optionally substituted heteroaryl group having 5 or 6 ring members, R^2 represents a cyclohexyl or cycloheptyl group optionally substituted by an alkyl group, R^3 represents a hydrogen atom and R^4 and R^5 , which may be identical or different, each represents a hydrogen atom or an alkyl group, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.

20 7. Compound of formula (I) according to any one of claims 1, 2, 5 and 6 wherein the alkyl group is a methyl group, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid or base.


25 8. Compound of formula (I) according to any one of claims 1 to 7 which is 2-[cyclohexyl(3-thienyl)methyl]-4-methyl-4,5-dihydro-1*H*-imidazole, the enantiomers, diastereoisomers and tautomers thereof, and also addition salts thereof with a pharmaceutically acceptable acid.

9. Process for the preparation of compounds of formula (I) according to claim 1, characterised in that there is used as starting material a compound of formula (II) :



wherein R^1 , R^2 and R^3 are as defined for formula (I),

- 5 which compounds of formula (II) are condensed with a diamine (III) :



wherein R^4 and R^5 are as defined for formula (I)

to yield, in the presence of an appropriate catalyst, compounds of formula (I),

- which may optionally be purified according to a conventional purification technique,
- which, if desired, are separated into their stereoisomers according to a conventional separation technique,
- which, if desired, are converted into additions salts with a pharmaceutically acceptable acid or base.

10. Pharmaceutical composition comprising as active ingredient at least one compound according to any one of claims 1 to 8, alone or in combination with one or more pharmaceutically acceptable, inert, non-toxic excipients or carriers.

11. Pharmaceutical composition according to claim 10 containing at least one active ingredient according to any one of claims 1 to 8 for use in the preparation of medicaments for the treatment of non-insulin-dependent type II diabetes, obesity,

type I diabetes, hyperlipidaemia, hypercholesterolaemia and cardiovascular complications thereof.

- 5 **12.** Pharmaceutical composition according to claim 10 containing at least one active ingredient according to any one of claims 1 to 8 for use in the preparation of medicaments for the treatment of type I and type II diabetes and cardiovascular complications thereof.
- 13.** Pharmaceutical composition according to claim 10 containing at least one active ingredient according to any one of claims 1 to 8 for use in the preparation of medicaments for the treatment of type I and type II diabetes.

**INPI NATIONAL INSTITUTE
FOR INDUSTRIAL PROPERTY**
**PATENT OF INVENTION
UTILITY CERTIFICATE**

Intellectual Property Code - Book VI

PATENTS DIVISION
 26 bis, rue de Saint Pétersbourg
75800 Paris Cedex 08

Tel.: 01 53 04 53 04

Fax: 01 42 93 59 30

DECLARATION OF INVENTORSHIP Page No. 1/3

(if the Applicant is not the inventor or not the only inventor)

This form is to be completed legibly in black ink

Your references for this file (<i>optional</i>)		70097	
NATIONAL REGISTRATION NO.		0213194	
TITLE OF THE INVENTION (maximum 200 characters or spaces)			
New imidazoline compounds, a process for their preparation and pharmaceutical compositions containing them			
APPLICANT(S):			
LES LABORATOIRES SERVIER 12, place de La Défense 92415 COURBEVOIE Cedex FRANCE			
DESIGNATE(S) AS INVENTOR(S) : (Indicate at the top right-hand side "Page No. 1/1". If there are more than three inventors, use an identical form and number each page indicating the total number of pages).			
Name		RAULT	
Forenames		Sylvain	
Address	Street	Route de Saint-Pierre sur Dives	
	Postal code and town	14370	MOULT (France)
Belonging company (<i>optional</i>)			
Name		KOPP	
Forenames		Marina	
Address	Street	33, rue Saint Martin	
	Postal code and town	14400	BAYEUX (France)
Belonging company (<i>optional</i>)			
Name		LANCELOT	
Forenames		Jean-Charles	
Address	Street	Tour en Bessin RN 13	
	Postal code and town	14400	LE BOURG (France)
Belonging company (<i>optional</i>)			
DATE AND SIGNATURE(S) OF THE APPLICANT(S) OR OF THE AUTHORISED AGENT (Name and position of signatory)		(signature)	
23 October 2002			
François RIVIERE, Patent Engineer			

**INPI NATIONAL INSTITUTE
FOR INDUSTRIAL PROPERTY**
**PATENT OF INVENTION
UTILITY CERTIFICATE**

Intellectual Property Code - Book VI

PATENTS DIVISION
 26 bis, rue de Saint Pétersbourg
75800 Paris Cedex 08

Tel.: 01 53 04 53 04

Fax: 01 42 93 59 30

DECLARATION OF INVENTORSHIP Page No. 2/3

(if the Applicant is not the inventor or not the only inventor)

This form is to be completed legibly in black ink

Your references for this file (optional)		70097	
NATIONAL REGISTRATION NO.		0213194	
TITLE OF THE INVENTION (maximum 200 characters or spaces) New imidazoline compounds, a process for their preparation and pharmaceutical compositions containing them			
APPLICANT(S): LES LABORATOIRES SERVIER 12, place de La Défense 92415 COURBEVOIE Cedex FRANCE			
DESIGNATE(S) AS INVENTOR(S) : (Indicate at the top right-hand side "Page No. 1/1". If there are more than three inventors, use an identical form and number each page indicating the total number of pages).			
Name		LEMAITRE	
Forenames		Stéphane	
Address	Street	29, rue des Bruyères	
	Postal code and town	27220	BOIS LE ROY (France)
Belonging company (optional)			
Name		CAIGNARD	
Forenames		Daniel-Henri	
Address	Street	22, avenue de la République	
	Postal code and town	78230	LE PECQ (France)
Belonging company (optional)			
Name		BIZOT-ESPIARD	
Forenames		Jean-Guy	
Address	Street	2, rue de la Convention	
	Postal code and town	75015	PARIS
Belonging company (optional)			
DATE AND SIGNATURE(S) OF THE APPLICANT(S) OR OF THE AUTHORISED AGENT (Name and position of signatory) 23 October 2002 François RIVIERE, Patent Engineer		(signature)	

received 08/11/02

**INPI NATIONAL INSTITUTE
FOR INDUSTRIAL PROPERTY**

**PATENT OF INVENTION
UTILITY CERTIFICATE**

Intellectual Property Code - Book VI

PATENTS DIVISION

26 bis, rue de Saint Pétersbourg
75800 Paris Cedex 08

Tel.: 01 53 04 53 04

Fax: 01 42 93 59 30

DECLARATION OF INVENTORSHIP Page No. 3/3

(if the Applicant is not the inventor or not the only inventor)

This form is to be completed legibly in black ink

Your references for this file (<i>optional</i>)		70097	
NATIONAL REGISTRATION NO.		0213194	
TITLE OF THE INVENTION (maximum 200 characters or spaces) New imidazoline compounds, a process for their preparation and pharmaceutical compositions containing them			
APPLICANT(S): LES LABORATOIRES SERVIER 12, place de La Défense 92415 COURBEVOIE Cedex FRANCE			
DESIGNATE(S) AS INVENTOR(S) : (Indicate at the top right-hand side "Page No. 1/1". If there are more than three inventors, use an identical form and number each page indicating the total number of pages).			
Name		RENARD	
Forenames		Pierre	
Address	Street	3, avenue du Parc	
	Postal code and town	78150	LE CHESNAY (France)
Belonging company (<i>optional</i>)			
Name			
Forenames			
Address	Street		
	Postal code and town		
Belonging company (<i>optional</i>)			
Name			
Forenames			
Address	Street		
	Postal code and town		
Belonging company (<i>optional</i>)			
DATE AND SIGNATURE(S) OF THE APPLICANT(S) OR OF THE AUTHORISED AGENT (Name and position of signatory) 23 October 2002 François RIVIERE, Patent Engineer		(signature)	

Law No. 78-17 of 6 January 1978 relating to information processing, data files and rights applies to the responses made on this form. It guarantees right of access to and correction of the data concerning you at the INPI.